

Fact Sheet:

White Peach Scale

Pseudaulacaspis pentagona

White Peach Scale (WPS) is considered a serious threat to the New Zealand kiwifruit industry. This pest has caused significant impact to the kiwifruit industry in Latina, Italy with fruit losses of 10 - 20% being recorded in some years. New Zealand has a suitable climate for this pest and could easily survive if established here.

Identification

WPS look similar to other armoured scale insects found in New Zealand. The adult female scale is immobile on the host plant and covered with a protective shell. It is dull white to yellowish in colour and oval in shape with an overall length of 2.0 - 2.5mm.



Figure 1. (From top) Adult females and larvae; numerous larvae; heavy infestation on 'Hongyang' kiwifruit in China (photo Garry Hill, Plant and Food Research)



Adult males emerge as tiny winged insects and begin building armour and develop a long white to

yellowish coloured shell. After moulting three more times they are orange and live for approximately 24 hours.

Distribution and Climate Range

WPS originated in eastern Asia and has global distribution including presence in USA, Europe, Australia and the Pacific Islands.

It is widely adapted to a range of climatic regions and could easily establish in New Zealand's conditions if it were introduced to a host here.




Figure 2. White Peach Scale distribution shown in yellow

Signs and Symptoms

WPS infects bark, fruit and leaves of plants. In severe cases it appears as white, cottony masses encrusting the bark of the tree. Scale insect feeding can result in early leaf and fruit drop. Trees which are heavily infested can become stunted and young plants can die very quickly after infestation.

Control

WPS is especially problematic when incursions occur in areas where natural predators are absent. Pheromone traps are widely used for detection in newly infested regions, especially in Europe. If an incursion does occur agricultural sprays such as those used against other armoured scale insects can be used. Costs of control are significant for countries facing incursions.

What should you do if you think you have seen this pest?

Phone: MPI - 0800809966 or KVH - 0800 665 825

Kiwifruit Vine Health Incorporated (KVH) makes no warranty or representation as to the accuracy or completeness of the information, photographs or other published material in this publication. KVH shall not be liable to any person for loss, injury or damages arising from a person's reliance on the published material. Published material authored by a person other than KVH reflects the view of the author and not necessarily the view of KVH. The published material may be subject to copyright and shall not be reproduced in any manner without first obtaining the permission of KVH.