

## Fact Sheet:

# Cherry Leaf Roll Virus

## Genus *Nepovirus*

Cherry leaf roll virus (CLRV) is one of two viruses known to induce severe symptoms in kiwifruit. CLRV was first identified in New Zealand on raspberries in 1978, and has been found to infect Hort16A kiwifruit.

### Identification

CLRV has symptoms similar to Psa-V and molecular (PCR) or serology (ELISA) testing is required to confirm its presence.

### Distribution and Climate Range



Figure 1. Known distribution of CLRV in all host species

CLRV has a worldwide distribution and a wide natural and experimental host range infecting members of more than 36 plant families. Hort16A is the only reported kiwifruit host of CLRV to date.

### Signs and Symptoms

Symptoms reported on Hort16A include leaf mottle and pale green spots that become necrotic. These necrotic lesions are larger and less angular than lesions induced by Psa-V. Cane dieback and severe stem cracking may also be seen. The characteristic “beak” of Hort16A fruit may be reduced or missing on CLRV-infected vines.



Figure 2. Absence of beak on CLRV-infected Hort16A fruit on right (Source: Plant and Food Research)



Figure 3. Severe stem cracking of CLRV infected vine (Source: Plant and Food Research)

### Control

CLRV is documented to be transmitted by seed, pollen, grafting and mechanical transmission. In kiwifruit, the virus has been observed to spread along rows as a result of pruning or girdling. Good orchard hygiene practices are required to control spread within an orchard, and nursery high health standards to prevent transmission through seed or nursery stock.

### To report vines displaying these symptoms:

Phone:

**MPI - 0800 80 99 66 or KVH - 0800 665 825**